

- Langeron O, Masso E, Huraux C, et al. Prediction of difficult mask ventilation. *Anesthesiology* 2000; **92**: 1229–1236.
- Levitan RM, Kinkle WC, Levin WJ, et al. Laryngeal view during laryngoscopy: a randomized trial comparing cricoid pressure, backward-upward-rightward pressure, and bimanual laryngoscopy. *Ann Emerg Med* 2006; **47**: 548–555.
- Neilipovitz DT, Crosby ET. No evidence for decreased incidence of aspiration after rapid sequence induction. *Can J Anaesthesiol* 2007; **54**: 748–764.
- Practice guidelines for management of the difficult airway: An updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology* 2003; **98**: 1269–1277.
- Ramachandran K, Kannan S. Laryngeal mask airway and the difficult airway. *Curr Opin Anaesthesiol* 2004; **17**: 491–493.
- Reed MJ, Dunn MJ, McKeown DW. Can an airway assessment score predict difficulty at intubation in the emergency department? *Emerg Med J* 2005; **22**: 99–102.
- Scrase I, Woollard M. Needle vs surgical cricothyroidotomy: a short cut to effective ventilation. *Anaesthesia* 2006; **61**: 921–923.
- Yentis SM. Predicting difficult intubation – worthwhile exercise or pointless ritual? *Anaesthesia* 2002; **57**: 105–109.
- Yildiz TS, Solak M, Toker K. The incidence and risk factors of difficult mask ventilation. *J Anesth* 2005; **19**: 7–11.